

01-03-00

Docket No. 58434-A/JPM/SHS

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Honorable Assistant Commissioner for Patents
Washington, D.C. 20231

December 29, 1999

S I R:

Transmitted herewith for filing are the specification and claims of the patent application of:

B. Jack Longley for
Inventor(s)METHODS FOR INHIBITING CUTANEOUS INFLAMMATION AND HYPERPIGMENTATION
Title of Invention

Also enclosed are:

☒ 9 sheet(s) of ☐ informal ☒ formal drawings.☒ Oath or declaration of Applicant(s). (unsigned)☒ A power of attorney (unsigned)☐ An assignment of the invention to☒ A Preliminary Amendment☐ A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT


	NUMBER FILED		NUMBER EXTRA*		RATE		FEE	
					SMALL ENTITY	OTHER ENTITY	SMALL ENTITY	OTHER ENTITY
Total Claims	27 -20	=	7	X	\$ 9.00	\$18.00	= \$ 63.00	\$ 0
Independent Claims	10 -3	=	7	X	\$39.00	\$78.00	= \$273.00	\$ 0
Multiple Dependent Claims Presented: Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>					\$130.00	\$260.00	= \$ 0	\$ 0
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE		\$ 345	\$ 690
					TOTAL FEE		\$681.00	\$ 0

Applicants: B. Jack Longley
Serial No.: Not Yet Known
Filed: December 29, 1999

Letter of Transmittal
Page 2

- ☒ A check in the amount of \$ 681.00 to cover the filing fee.
- ☐ Please charge Deposit Account No. _____ in the amount of \$ _____.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:
- ☒ Filing fees under 37 C.F.R. §1.16.
- ☒ Patent application processing fees under 37 C.F.R. §1.17.
- ☐ The issue fee set in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).
- ☒ Three copies of this sheet are enclosed.
- ☐ A certified copy of previously filed foreign application No. _____ filed in _____ on _____ Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.
- ☒ Other (identify) Express Mail Certificate of Mailing bearing label No. EM 262 551 295 US, dated December 29, 1999; 1 loose set of formal drawings

Respectfully submitted,



John P. White
Registration No. 28,678
Attorney for Applicant(s)
Cooper & Dunham, LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: B. Jack Longley

Serial No.: Not Yet Known (Continuation-in-part of
U.S. Serial No. 09/306,143,
filed May 6, 1999)

Filed : December 29, 1999

For : METHODS FOR INHIBITING CUTANEOUS INFLAMMATION
AND HYPERPIGMENTATION

1185 Avenue of the Americas
New York, New York 10036
December 29, 1999

Assistant Commissioner for Patents
Washington, D.C. 20231
Attn: Box Patent Application


SIR:

**EXPRESS MAIL CERTIFICATE OF
MAILING FOR ABOVE-IDENTIFIED APPLICATION**


"Express Mail" mailing label number: EM 262 551 295 US

Date of Deposit: December 29, 1999

I hereby certify that this paper or fee is being deposited with
the United States Postal Service "Express Mail Post Office to
Addressee" service under 37 C.F.R. §1.10 on the date indicated
above and is addressed to the Assistant Commissioner for Patents
Box Patent Applications, Washington, D.C. 20231.


Printed Name: John P. White

Respectfully submitted,


John P. White
Registration No. 28,678
Attorney for Applicant(s)
Cooper & Dunham, LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: B. Jack Longley
Serial No.: Not Yet Known (Continuation-in-part of
U.S. Serial No. 09/306,143,
filed May 6, 1999)
Filed : December 29, 1999
For : METHODS FOR INHIBITING CUTANEOUS INFLAMMATION
AND HYPERPIGMENTATION

1185 Avenue of the Americas
New York, New York 10036
December 29, 1999

Assistant Commissioner for Patents
Washington, D.C. 20231
Attn: Box Patent Application

SIR:

PRELIMINARY AMENDMENT

Please amend the subject application as follows:

In the claims:

Please cancel claims 10-11, 20-22, 27, 32 and 35-38 without prejudice to applicants' right to pursue the subject matter of these claims in a later-filed application. Please amend claims 12-19, 23-26, 28-31 and 33-34 under the provisions of 37 C.F.R. §1.121(b) by deleting the bracketed material and inserting the underlined material as follows.

- 12. (Amended) The method of [any one of claims 1-9] claim 1, which comprises inhibiting the kinase enzymatic reaction of kit protein.--
- 13. (Amended) The method of [any one of claims 1-9] claim 1, which comprises inhibiting chymase, elastase or other SCF cleaving enzymes.--
- 14. (Amended) The method of [any one of claims 1-9] claim 1, which comprises inhibiting ligand binding

Applicants: B. Jack Longley
Serial No.: Not Yet Known
Filed : December 29, 1999
Page 2

with an antibody, peptide, or nonpeptide chemical.--

--15. (Amended) The method of [any one of claims 1-9] claim 1, which comprises inhibiting kit dimerization with an antibody, peptide, or nonpeptide chemical.--

--16. (Amended) The method of [any one of claims 1-9] claim 1, wherein downstream signaling of the kit activation pathway is inhibited by blocking substrate association with kit kinase domain.--

--17. (Amended) The method of [any one of claims 1-9] claim 1, wherein downstream signaling of the kit activation pathway is inhibited by blocking enzymatic function in the downstream signaling pathway.--

--18. (Amended) The method of [any one of claims 1-9] claim 1, wherein downstream signaling of the kit activation pathway is inhibited by blocking binding of molecules in the downstream signaling pathway.--

--19. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound is an antibody or portion thereof.--

--23. (Amended) The method of claim [22] 19, wherein the [anti-kit] antibody is ACK2.--

--24. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound comprises a Fab

Applicants: B. Jack Longley
Serial No.: Not Yet Known
Filed : December 29, 1999
Page 3

fragment of an anti-kit antibody.--

--25. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound comprises the variable domain of an anti-kit antibody.--

--26. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound comprises one or more CDR portions of an anti-kit antibody.--

--28. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound comprises a peptide, peptidomimetic, a nucleic acid, or an organic compound with a molecular weight less than 500 Daltons.--

--29. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound is sSCF, sKIT ligand or a fragment thereof.--

--30. (Amended) The method of [any one of claims 1-9] claim 1, wherein the compound is sKIT or a fragment thereof.--

--31. (Amended) The method of [any one of claims 1-9] claim 1, wherein the subject is a mammal.--

--33. (Amended) The method of [any one of claims 1-9] claim 1, wherein the administration is intralesional, intraperitoneal, intramuscular, subcutaneous, intravenous, liposome mediated delivery, transmucosal, intestinal, topical, nasal, oral, anal, ocular or otic delivery.

Applicants: B. Jack Longley
Serial No.: Not Yet Known
Filed : December 29, 1999
Page 4


Remarks:

Claims 1-38 were pending in the subject application. Applicants have hereinabove canceled claims 10-11, 20-22, 27, 32 and 35-38 without prejudice to their right to pursue the subject matter of these claims in a later-filed application and amended claims 12-19, 23-26, 28-31 and 33-34. Claims 1-9, 12-19, 23-26, 28-31 and 33-34 involve no issue of new matter and entry of this amendment is respectfully requested.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicants' undersigned attorney invites the Examiner to telephone him at the number provided below.

No fee, in addition to the enclosed filing fee of \$681.00, is deemed necessary in connection with the filing of this Preliminary Amendment. However, if any additional fee is required, authorization is hereby given to charge the amount of any such fee to Deposit Account No. 03-3125.

Respectfully submitted,



John P. White
Registration No. 28,678
Attorney for Applicant(s)
Cooper & Dunham, LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

**Application
for
United States Letters Patent**

To all whom it may concern:

Be it known that B. Jack Longley

have invented certain new and useful improvements in

METHODS FOR INHIBITING CUTANEOUS INFLAMMATION AND HYPERPIGMENTATION

of which the following is a full, clear and exact description.

**METHODS FOR INHIBITING CUTANEOUS
INFLAMMATION AND HYPERPIGMENTATION**

5 This application is a continuation-in-part of U.S. Serial
No. 09/306,143, filed May 6, 1999, the contents of which
are hereby incorporated by reference into this
application.

10 The invention described herein was made with Government
support under grant numbers 1 R29 AR 40514-01A1, 5 P30
041942 and 1-RO1-AR43356-01A2 from the National
Institutes of Health. Accordingly, the United States
Government has certain rights in this invention.

15 Throughout this application, various publications are
referenced by arabic numerals within parentheses.
Disclosures of these publications in their entireties are
hereby incorporated by reference into this application to
more fully describe the state of the art to which this
20 invention pertains. Full bibliographic citations for
these references may be found immediately preceding the
claims.

Background of the Invention

25 The use of murine models to investigate human cutaneous
oncology, immunology and keratinocyte biology is
advantageous over the use of human skin for obvious
reasons. However, substantial differences exist between
human skin and murine skin. In human skin, Stem Cell
30 Factor is produced by epidermal keratinocytes after
birth, unlike in normal murine skin. The result of this,
among other things, is that melanocytes are present in
the interadnexal epidermis in human skin. In contrast,
melanocytes in adult murine skin are generally confined
35 to hair follicles, with the exception of rare epidermal
melanocytes found in the ears, footpads, and tail (1).
A few dermal melanocytes may also be found in mice,

mostly in the ears. These differences have compromised the use of the mice as a model system for investigation of human cutaneous biology.

5 It has been discovered that melanocyte migration and development, as well as the survival of melanocytes and mast cells, are dependent on expression of the kit protein, a receptor tyrosine kinase encoded by the c-kit proto-oncogene (2-6). The ligand for kit, known as stem
10 cell factor (SCF) (also called mast cell growth factor, steel factor, and kit ligand) may be produced locally in human skin by epidermal keratinocytes, fibroblasts, and endothelial cells (7-8). However, definitive studies of SCF production in murine skin have not been reported.
15 Transgenic studies using the SCF gene promoter region and beta-galactosidase as a reporter gene suggest that, unlike in human skin, postnatal murine cutaneous SCF expression is limited to the dermis and hair follicles, and not found in epidermal keratinocytes (9). The
20 difference in SCF expression between human and murine epidermis could explain the difference in melanocyte distribution and other biological phenomena in these two species.

25 SCF may be produced in two isoforms by alternate splicing of exon 6. One isoform lacks exon 6 encoded sequences and exists predominantly as a membrane-bound molecule. The other isoform contains exon 6 encoded sequences which include a protease sensitive site (10-19). Cleavage at
30 the protease sensitive site causes the release of a soluble, bioactive form of SCF. The membrane-bound and soluble forms of SCF have differential effects on melanocyte precursor dispersal and survival (20) and exogenous soluble SCF may produce cutaneous mast cell
35 hyperplasia and cutaneous hyperpigmentation (21-23). In addition, local high concentrations of soluble SCF have been found in lesions of human cutaneous mastocytosis, a

disease characterized by dermal accumulations of mast cells and increased epidermal melanin (7, 8, 24) and in spongiotic dermatitis, a common inflammatory condition of human skin (our unpublished data).

5

Summary of the Invention

This invention provides a method of preventing or treating in a subject contact dermatitis which comprises
10 administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat contact dermatitis so as to thereby prevent or treat contact dermatitis in the subject.

15

This invention provides a method of preventing or treating in a subject hyperpigmentation which comprises
20 administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat hyperpigmentation so as to thereby prevent or treat hyperpigmentation in the subject.

This invention provides a method of preventing or
25 treating in a subject asthma which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat asthma so as to thereby prevent or treat asthma in the subject.

30

This invention provides a method of preventing or treating in a subject cutaneous inflammation which comprises administering to the subject an amount of a
35 compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat cutaneous inflammation so as to thereby prevent or treat cutaneous inflammation in the subject.

5 This invention provides a method of preventing or treating in a subject anaphylaxis and bronchospasm which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat anaphylaxis and bronchospasm so as to thereby prevent or treat anaphylaxis and bronchospasm in the subject.

10 This invention provides a method of preventing or treating in a subject mastocytosis which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat mastocytosis so as to thereby prevent or treat mastocytosis in the subject.

15 This invention provides a method of preventing or treating in a subject urticaria which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat urticaria so as to thereby prevent or treat urticaria in the subject.

20 This invention provides a method of preventing or treating in a subject hypersensitivity reactions which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat urticaria so as to thereby prevent or treat urticaria in the subject.

25 This invention provides a method of preventing or treating in a subject a tumor which expresses activated kit which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat a tumor which expresses activated kit so as to thereby prevent or treat a tumor which expresses activated kit in

30

35

the subject.

5 The invention provides a method of providing
contraception to a subject which comprises administering
to the subject an amount of a compound capable of
inhibiting the stem cell factor signaling pathway
effective to prevent conception so as to thereby provide
contraception to the subject.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2

Brief Description of the Figures

Figure 1:

Transgene design. Both transgenes used the human keratin
14 promoter and polyadenylation sequences. Transgene one
5 included a rabbit b-globin intron, and transgene two
included human growth hormone sequences to provide for
stability. Neither the beta globin intron nor the human
growth hormone sequences produce protein products.

Figure 2:

Increased mast cells in mice expressing epidermal
membrane and soluble SCF (transgene one). (a) Numerous
mast cells are demonstrated in the superficial dermis of
15 body wall skin of newborn mice bearing transgene one
(membrane/soluble SCF), using an immunoperoxidase/alcian
blue technique which stain mast cell granules
metachromatically purple. Note the apposition of mast
cells (arrowheads) to basilar keratinocytes, the source
20 of SCF. Immunoperoxidase with an anti-S100 antibody in
this preparation also demonstrates melanocytes as brown
staining cells in the epidermal basilar layers of
epidermis and follicles (white arrows). Sebocytes are
seen as large, round, lightly S-100(+) cells in the
25 follicular epithelium. Melanin pigment is stained black
in this preparation. (b) Immunofluorescence with anti-kit
antibodies highlights kit expressing dermal mast cells
(arrowheads) in body wall skin of newborn (transgene one
membrane/soluble SCF) mouse. (c) Anti-kit antibody
30 immunofluorescence shows mast cells crowded in the
papillary dermis and extending into the upper reticular
dermis and body wall skin of 21 day old, transgene one
positive mouse, MC, confluent mast cells; arrowheads,
individual and small clusters of mast cells; E,
35 epidermis; F, follicles; K, keratin layer. (d)
Hematoxylin and eosin-stained sections show mast cells
filling the superficial corium in section of tongue from

a 21 day old, transgene one positive mouse. The lack of abundant melanocytes and melanophages in this anatomic site allows easy visualization of the mast cells. This histologic picture is identical to that seen in human cutaneous mastocytosis. (e) Alcian blue stained serial section of tongue shows metachromatic granules in mast cells of 21 day old, transgene one positive mouse.

Figure 3:

Electron microscopy confirms the presence of melanocytes and mast cells in transgenic mice. (a) Transgene one mouse with membrane/soluble epidermal SCF has numerous dermal mast cells (arrowheads) as well as dermal melanocytes (arrows). Asterisks show the boundary of the dermis and hair follicle. Higher magnification images of mast cell and melanocyte are shown in b and c, respectively. Original magnifications: (a) 2,750, (b) 9,000, (c) 11,750.

Figure 4:

Transgenic phenotypes are stable across a wide range of gene expression levels. This figure compares the transgene copy number determined by PCR, with SCF mRNA expression as determined by RNase protection assay, in lines from different founders. The relative density of SCF bands was determined by dividing the mean density of the SCF band by the density of a SCF band derived from an identical aliquot of RNA. Probe templates were 384 bases in length for SCF (40 base pairs of promoter sequence and 342 bases complimentary to nucleotides 814-1156 of murine SCF mRNA (5). A beta-actin probe was used as a control, and to allow standardization between RNA preparations from different mice. The beta-actin probe length was 310 bases, 227 bases of which are complementary to murine beta-actin mRNA. The probe was purchased from Ambion (pTR1-beta-actin-mouse anti-sense control template). Note the differences between TG2 (4x, 5x, 10x) and TG1

(6x).

Figure 5:

Epidermal SCF causes hyperpigmentation of murine skin.

5 (a) Newborn mouse expressing membrane/ soluble SCF
(transgene one, left) shows obvious hyperpigmentation
compared to non-transgenic littermate (right). (b)
Transgene two positive mouse overexpressing
10 membrane-bound epidermal SCF shows a similar phenotype
with generalized hyperpigmentation which is most
discernible in the ventral and hairless areas, and which
is maintained in adult life. Three week old transgenic
(left) and non-transgenic littermate (right)

15 Figure 6:

Intraepidermal melanocytes are increased in transgenic
mice. (a) Tail skin section from 21 day old mouse

expressing epidermal membrane-bound SCF (transgene two)
shows mild epidermal hyperplasia and a markedly increased
20 number of melanocytes, identified as cells surrounded by
clear halos, mostly at the dermal-epidermal junction.
These mice also show extensive black epidermal melanin
pigment (400x). (b) Note the lack of both basilar
melanocytes and epidermal pigment in the skin of the
25 transgene (-) littermate control mouse (C57 black 6
(400x)). (c) Epidermal melanocytes express kit protein.
Immunofluorescence staining with anti-kit antibody and
Texas Red labeled secondary antibody demonstrates
confluent dendritic cells in the epidermal basalar layer
30 of mice expressing membrane-bound SCF (transgene two
arrows). These cells correspond to the S-100 protein (+)
basilar dendritic cells seen in Figure 2a. Note two
strongly kit positive solitary mast cells in the dermis
(arrowheads, 400x). Light staining of dendritic
35 melanocytes can also be seen in the epidermis of
transgene one positive mice (please see figure 2b).

Figure 7:

Electron microscopy confirms the presence of epidermal melanocytes in both types of transgenic mice. (a) Electron microscopy shows numerous keratinocytes containing phagocytized melanin granules in the interadnexal epidermis of mice expressing membrane-bound epidermal SCF (3500x). b. Epidermal melanosomes, some marked with large arrows, are present in both keratinocytes and melanocytes. Pre melanosomes, marked with the open arrows, demonstrate the presence of a melanocyte. Note keratinocyte hemidesmosomes (small arrows) which confirm the location of the melanocyte within the epidermis (16, 320x).

Figure 8:

Allergic ear swelling is significantly increased in SCF transgenic animals, and is reduced by blocking the SCF receptor with the ACK2 monoclonal antibody. All transgenic mice show increased ear swelling in response to allergic contactants compared to non-transgenic animals ($p \leq 0.0001$), showing that SCF contributes to dermatitis. Similar results are seen with irritant contactants (data not shown). The ear swelling is specifically decreased by the ACK2 monoclonal antibody which blocks the SCF receptor ($p \leq 0.05$) confirming that epidermal SCF plays an active role in cutaneous inflammation.

Figure 9:

(a) An unpublished immunoperoxidase study of inflamed human skin with an anti-human SCF monoclonal anti-body shows soluble epidermal SCF in spongiotic (eczematous) dermatitis, here demonstrated in an epidermal spongiotic vesicle. Human spongiotic dermatitis may be associated with hyperpigmentation. Soluble epidermal SCF is not detected in normal skin with this technique (7), suggesting that epidermal SCF is released in cutaneous

inflammatory states. (b) Irritant dermatitis induced in transgenic mice by Croton oil is characterized histologically by spongiotic (eczematous) dermatitis. The murine dermatitis is shown in a hematoxylin and eosin stained slide because no satisfactory antibody against murine SCF is available to allow demonstration of soluble SCF by immunoperoxidases. However, these results combined with the ACK2 blocking studies shown above implicate epidermal SCF in this phenomenon.

5

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209

Detailed Description of the Invention

5 This invention provides a method of preventing or treating in a subject contact dermatitis which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat contact dermatitis so as to thereby prevent or treat contact dermatitis in the subject.

10

15 This invention provides a method of preventing or treating in a subject hyperpigmentation which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat hyperpigmentation so as to thereby prevent or treat hyperpigmentation in the subject.

20 This invention also provides a method of preventing or treating spongiotic dermatitis which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat spongiotic dermatitis so as to thereby prevent or treat spongiotic dermatitis in the subject. As used herein, "spongiotic dermatitis" includes but is not limited to contact dermatitis.

25

30 This invention provides a method of preventing or treating in a subject asthma which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat asthma so as to thereby prevent or treat asthma in the subject.

35

This invention provides a method of preventing or treating in a subject cutaneous inflammation which

comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat cutaneous inflammation so as to thereby prevent or treat cutaneous inflammation in the subject.

This invention provides a method of preventing or treating in a subject anaphylaxis and bronchospasm which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat anaphylaxis and bronchospasm so as to thereby prevent or treat anaphylaxis and bronchospasm in the subject.

This invention provides a method of preventing or treating in a subject mastocytosis which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat mastocytosis so as to thereby prevent or treat mastocytosis in the subject.

This invention provides a method of preventing or treating in a subject urticaria which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat urticaria so as to thereby prevent or treat urticaria in the subject.

This invention provides a method of preventing or treating in a subject hypersensitivity reactions which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat hypersensitivity reactions so as to thereby prevent or treat hypersensitivity reactions in the subject.

5 This invention provides a method of preventing or treating in a subject a tumor which expresses activated kit which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat a tumor which expresses activated kit so as to thereby prevent or treat a tumor which expresses activated kit in the subject.

10 In one embodiment, the tumor is a mast cell tumor. In another embodiment, the tumor is a gastrointestinal stromal tumor. In another embodiment, the tumor is a germ cell tumor.

15 In one embodiment, the above method comprises inhibiting the kinase enzymatic reaction of kit protein.

In one embodiment, the above method comprises inhibiting chymase, elastase or other SCF cleaving enzymes.

20 In one embodiment, the above method comprises inhibiting ligand binding with an antibody, peptide, or nonpeptide chemical.

25 In one embodiment, the above method comprises inhibiting receptor dimerization with an antibody, peptide, or nonpeptide chemical.

30 In one embodiment of the above method, downstream signaling of the kit activation pathway is inhibited by blocking substrate association with the kit kinase domain.

35 In one embodiment of the above method, downstream signaling of the kit activation pathway is inhibited by blocking enzymatic function in the downstream signaling pathway.

In one embodiment of the above method, downstream signaling of the kit activation pathway is inhibited by blocking binding of molecules in the downstream signaling pathway.

5

In one embodiment of the above method, the compound is an antibody or portion thereof. In one embodiment, the antibody is a monoclonal antibody. In one embodiment, the monoclonal antibody is a human, humanized or a chimeric antibody. In one embodiment, the monoclonal antibody is an anti-kit antibody. In one embodiment, the anti-kit antibody is ACK2.

10

This invention provides the above method, wherein the compound comprises a Fab fragment of an anti-kit antibody.

15

This invention provides the above method, wherein the compound comprises the variable domain of an anti-kit antibody.

20

This invention provides the above method, wherein the compound comprises one or more CDR portions of an anti-kit antibody.

25

This invention provides the above method, wherein the antibody is selected from the group consisting of IgA, IgD, IgE, IgG and IgM.

30

This invention provides the above method, wherein the compound comprises a peptide, peptidomimetic, a nucleic acid, or an organic compound with a molecular weight less than 500 Daltons.

35

This invention provides the above method, wherein the compound is sSCF, sKIT ligand or a fragment thereof. As used herein "sSCF" can also mean "sKIT ligand."

This invention provides the above method, wherein the compound is SKIT or a fragment thereof.

5 This invention provides the above method, wherein the subject is a mammal. The subject of the above methods includes but is not limited to a mammal. The subject may be a mammal or non-mammal. The subject may be a human, a primate, an equine subject, an opine subject, an avian subject, a bovine subject, a porcine, a canine, a feline
10 or a murine subject. In another embodiment, the subject is a vertebrate. In a preferred embodiment, the mammal is a human being.

15 This invention provides the above method, wherein the administration is intralesional, intraperitoneal, intramuscular, subcutaneous, intravenous, liposome mediated delivery, transmucosal, intestinal, topical, nasal, oral, anal, ocular or otic delivery.

20 This invention provides a method of providing contraception to a subject which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent conception so as to thereby provide
25 contraception to the subject. In one embodiment, the subject is a male subject. In another embodiment, the subject is a female subject.

30 This invention provides the above method which comprises inhibiting the kinase enzymatic reaction of kit protein.

This invention provides the above method comprises inhibiting chymase, elastase or other SCF cleaving enzymes.

35

One skilled in the art can employ various methods for determining whether a compound inhibits the stem cell

signaling pathway. One of these methods comprises:

- a) immobilizing kit protein on a solid matrix;
- b) contacting the immobilized kit protein with the compound being tested and a predetermined amount of SCF under conditions permitting binding of kit protein and SCF in the absence of the compound;
- c) removing any unbound compound and any unbound SCF;
- d) measuring the amount of SCF which is bound to the immobilized kit protein;
- e) comparing the amount measured in step (d) with the amount measured in the absence of the compound, a decrease in the amount of SCF bound to the kit protein in the presence of the compound indicating that the compound inhibits binding of SCF to kit protein, thereby indicating that the compound inhibits the stem cell factor signaling pathway.

Another method includes detecting in vitro phosphorylation of kit comprising the following steps:

1. Cells expressing KIT, either naturally occurring as in the C2, BR, P815, or HMC1 lines, or as the result of DNA transfection, are grown in vitro in the absence of exogenous SCF.
2. The cells are treated with the compound or substance. In the case of cells expressing wild type (non-mutated and not constitutively activated) KIT, treatment with the substance or compound may be followed by treatment with SCF. Controls include cells treated with compound or substance, and not exposed to SCF.
3. The cells are lysed and KIT is immunoprecipitated, electrophoresed, blotted with anti-phosphotyrosine antibody, and the

antibody detected by chemiluminescence or radioactive labeling and autoradiography, thereby determining the level of KIT phosphorylation on tyrosine.

5

4. The level of KIT phosphorylation on tyrosine in cells not exposed to the compound or substance is compared to KIT phosphorylation on tyrosine in cells exposed to the compound or substance. A decrease in KIT phosphorylation on tyrosine in the treated cells indicates that the compound or substance inhibits the SCF signaling pathway.

10

A cell proliferation and viability method comprises the following steps:

15

1. Cells expressing KIT, and depending on KIT activation for survival, are grown in the presence of SCF if they express wild type KIT or in the absence of SCF if they express mutated and constitutively activated KIT.
2. The cells are grown in replicate tissue culture wells, in the presence or absence of the compound or substance to be tested, and the number of viable and non-viable cells per well is determined daily by counting a sample of cells in a hemocytometer. The number of viable and non-viable cells are determined by the trypan blue exclusion method.
3. A decrease in cell growth, as determined by the presence of fewer viable cells in the wells treated with the compound or substance compared to the cells in the wells not so treated, indicates that the compound or substance interferes with the KIT SCF signaling pathway.
4. Alternatively, cell growth may be determined by measuring cellular incorporation of labeled

20

25

30

35

substances such as tritium labeled thymidine.

5 The present invention provides a method of identifying a composition, a compound or a procedure which can produce a skin response in a subject, comprising: a) administering said composition or compound, or applying
10 said procedure to the transgenic mice which express endogenous epidermal stem cell factor, and b) analyzing the contacted skin for response.

15 In one embodiment of the method, the composition or compound can be administered orally or by injection.

In another embodiment of the method, the composition or compound can be administered topically by contacting the composition or compound with the skin of the transgenic mice.

20 In another embodiment of the method, the procedure is not previously known.

25 In another embodiment of the method, the procedure is identified by the method.

In another embodiment of the method, the procedure is DNA vaccination.

30 In this invention, the skin response may be induced. This skin response includes but is not limited to inflammation, tanning, melanoma, carcinoma or hyperpigmentation.

35 In another embodiment of the method, the composition may be cosmetics, medications or skin care products.

In another embodiment of the method, the composition or compound is not previously known.

5 In yet another embodiment of the method, the composition or compound is identified by the method.

10 In a further embodiment of the method, a mixture is produced for producing a skin response comprising an effective amount of the composition or compound identified by the method and a suitable carrier.

15 The present invention also provides a method of identifying a composition, a compound, or a procedure which can reduce or treat skin response in a subject, comprising: a) administering said composition or compound, or applying said procedure to the transgenic mice which express endogenous epidermal stem cell factor and which had been induced to produce a skin response and b) analyzing the skin of said transgenic mice to
20 determine the reduction of skin response, wherein the reduction of skin response indicates that the composition, compound, or procedure can reduce skin response.

25 In one embodiment of the method, the composition or compound can be administered orally or by injection.

30 In another embodiment of the method, the composition or compound can be administered topically by contacting the composition or compound with the skin of the transgenic mice.

In another embodiment of the method, the procedure is not previously known.

35 In another embodiment of the method, the procedure is identified by the method.

In another embodiment of the method, the procedure is DNA vaccination.

5 In another embodiment of the method, the composition or compound is not previously known.

In another embodiment of the method, the composition or compound is identified by the method.

10 In another embodiment of the method, a mixture is produced for reducing skin response comprising an effective amount of the composition or compound identified by the method and a suitable carrier.

15 In another embodiment of the method, the skin response is inflammation, tanning, skin carcinoma, melanoma or hyperpigmentation.

20 In another embodiment of the method, the hyperpigmentation is natural occurring hyperpigmentation or post inflammatory hyperpigmentation.

25 In another embodiment of the method, the inflammation is associated with human hyperpigmentation, or human hypopigmentation.

In another embodiment of the method, the subject is a mouse or a human-being.

30 In another embodiment of the method, the epidermal stem cell factor transgene encodes either a membrane bound epidermal stem cell factor or a membrane/soluble epidermal stem cell factor.

35 In another embodiment of the method, the epidermal stem cell factor transgene encodes a membrane or soluble epidermal stem cell factor.

In another embodiment of the method, the epidermal stem cell factor transgene is cloned into a construct containing a human cytokeratin 14 promotor.

5 In another embodiment of the method, the human cytokeratin 14 promotor causes the expression of the stem cell factor transgene in murine skin of the basal layers of the interadnexal epidermis and the follicular epithelium.

10

In another embodiment of the method, the skin response of the transgenic mice can be induced by applying an irritant or an allergic dermatitis inducing agent to said skin.

15

In another embodiment of the method, the irritant is croton oil or dinitrofluorobenzene.

20

In another embodiment of the method, the croton oil or dinitrofluorobenzene are applied to the ear or the abdominal skin of the transgenic mice; wherein the abdominal skin is either hairless or shaved.

25

In another embodiment of the method, the croton oil is used at a concentration of 0.2 percent.

30

In another embodiment of the method, the dinitrofluorobenzene is used at a concentration of 0.5 percent in a 4:1 mixture of acetone and olive oil.

In another embodiment of the method, the reduction or treatment of hyperpigmentation is determined by electron microscopic analysis.

35 In another embodiment of the method, the compound is an epidermal stem cell factor inhibitor.

In yet another embodiment of the method, the stem cell factor inhibitor is a monoclonal antibody.

5 In a further embodiment of the method, the monoclonal antibody is ACK2.

10 The present invention further provides a method of identifying a composition, a compound or a procedure which can reduce radiation damage to the skin of a subject, comprising: a) administering said composition or compound, or applying said procedure to the transgenic mice which express endogenous epidermal stem cell factor, b) subjecting the skin of said transgenic mice and the skin of the control transgenic mice to radiation, and c) 15 analyzing the effects of said composition, compound, or procedure on reducing skin radiation damages.

20 In one embodiment of the method, the composition or compound can be administered orally or by injection.

25 In another embodiment of the method, the composition or compound can be administered topically by contacting the composition or compound with the skin of the transgenic mice.

30 In another embodiment of the method, the procedure is not previously known.

35 In another embodiment of the method, the procedure is identified by the method.

In another embodiment of the method, the procedure is DNA vaccination.

In another embodiment of the method, the composition or compound is not previously known.

In another embodiment of the method, the composition or compound is identified by the method.

5 In another embodiment of the method, a mixture is produced for reducing skin radiation damages comprising an effective amount of the composition or compound identified by the method and a suitable carrier.

10 In yet another embodiment of the method, the radiation is ultra-violet light.

15 In a further embodiment of the method, the radiation damage is tanning, carcinogenesis, photo-aging, photo-damage or the development of melanoma.

20 The present invention also provides a pharmaceutical composition for treating human skin diseases, comprising (a) a compound that can treat skin diseases of the transgenic mice which express endogenous epidermal stem cell factor, and (b) a suitable carrier, wherein the compound specifically targets the epidermal stem cell factor or its receptor.

25 In one embodiment of the pharmaceutical composition, the compound is ACK2.

30 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

35

Experimental Details

Transgene construction:

Two murine SCF cDNAs were cloned into constructs containing the human cytokeratin 14 upstream region (27) (Figure 1). This promoter causes expression in the skin limited to the basal layers of the interadnexal epidermis and the follicular epithelium. The cDNAs were both full length clones, containing exon 6 encoded sequences. One cDNA (transgene one) was unmodified and therefore could produce a membrane-bound protein with the exon 6 encoded protease sensitive site, from which a soluble, bioactive form of SCF could be efficiently generated (10, 11, 28). The product of this transgene will be referred to as membrane/soluble SCF. The second cDNA (transgene two) had been previously modified by site directed mutagenesis, deleting the primary high efficiency cleavage site (between amino acids 164 and 165) and an alternate exon 7 encoded low efficiency cleavage site (found in murine SCF between amino acids 180 and 181). The SCF produced by this transgene therefore exists predominantly as a membrane-bound molecule (membrane SCF) (29). Both cDNAs have been previously shown to produce biologically active SCF (29, 30).

Generation and analysis of transgenic animals:

Two $\mu\text{g/ml}$ transgenic DNA, in 10 mM Tris (pH 7.5), 0.1 mM EDTA was injected into fertilized oocytes collected from pseudopregnant mice as described (31). At birth, most transgene expressing mice could be identified by distinctive pigmentary phenotypes, as described in the Results section. Integration of transgenes was verified by polymerase chain reaction (PCR) of genomic DNA with transgene-specific primers and copy number estimated by Southern blotting of PCR products, followed by autoradiography and densitometry. Skin specific expression of transgene messenger RNA was confirmed by northern blotting and by reverse transcription-polymerase chain reaction with transgene specific primers using RNA

5 extracted from representative animals. Transgene
expression was quantitated by RPA II Ribonuclease
Protection Assay Kit (Ambion, Austin, TX, USA) according
to the manufacturer's directions. Briefly, total RNA
10 extracted from mouse skin was hybridized with digoxigenin
labeled single stranded RNA probes for twenty three hours
at 42°C, digested with RNase A and RNase T1,
electrophoresed through 5% polyacrylamide/7 Molar urea,
protected fragments were transferred to Hy⁺ membrane
15 (Boerheringer-Mannheim, Indianapolis, IN, USA), bands
detected by chemiluminescences, and band density
determined by densitometry. Preliminary studies of RNA
preparations from each transgenic line were performed to
measure Beta-actin, and the amounts of RNA for SCF mRNA
20 determinations adjusted for comparison. RNA was also
used with reverse transcription and the polymerase chain
reaction for direct ampilmer sequencing of c-kit mRNA
sequences in regions which could contain known activating
mutations, as previously described (25). The primers
used were 5' CAAATC/GCATCCC/TCACACCCTGTTCAC and 5'
25 CCATAAGCAGTTGCCTCAAC which bind to nucleotides 1568→1593
and 1854→1835 and 5' TGTATTACAGAGACTTGGC and 5'
AAAATCCCATAGGACCAGAC binding to nucleotides 2384→2403 and
2595→2576. These regions contain the codons with both of
the activating mutations, codon 559 and codon 814,
respectively which have been described in human
mastocytosis and in a murine mast cell line (5, 26).

30 Transgene one, containing the full-length unmodified SCF
cDNA (membrane/soluble SCF), was injected into 100 F1
oocytes (C57 BL6 x SLJ), which were implanted into six
host mothers, resulting in four independent
hyperpigmented mice, all of which were positive for the
transgene, and 40 other littermates which were pigmentary
35 phenotype negative and transgene negative by PCR.

Oocytes for transgene two (membrane SCF) were F1

(C57BL/6J female x SLJ/J male), and the offspring could be black, agouti, or white. Injection of 40 embryos and implantation into six host mothers generated 48 pups, 21 of which were positive for integration by PCR. Of the 25 founder mice identified by PCR with the transgene specific primers, 3 were black, 13 were agouti, and 9 were white. Five PCR positive mice (3 agouti and 2 black) showed a clearly identifiable pigmentary phenotype. Given the inability of white mice to produce normal cutaneous pigment, it is possible that there were also white founders that expressed the transgene without the production of an obvious change in pigment. Backcrossing of phenotype positive, black and agouti founders to C57 BL/6 mice produced uniform pigmentary changes, described in the Results section.

Histology, immunohistochemistry, and electron microscopy:
Tissues from transgenic and littermate mice were fixed in formalin and embedded in paraffin or polyester wax, sectioned, and stained with hematoxylin and eosin, azure blue, alcian blue, or Giemsa's stain according to standard techniques (31-33). Immunofluorescence studies were performed on polyester wax embedded sections or frozen sections, also using standard techniques. Antibodies included anti-S100 (rabbit anti-cow S100, pre-diluted, Dako, Carpinteria, CA), and the ACK2 and ACK4 monoclonals (rat anti-mouse c-kit (34), at 20 μ g/ml). Controls included omission of the primary antibody or the use of isotype matched monoclonal antibodies of irrelevant specificity. Electron microscopy was done as previously described (35).

Inflammation Inducement and Treatment
We used Croton Oil and dinitrofluorobenzene (DNFB), respectively to reduce irritant and allergic contact dermatitis, respectively, in HK14-SCF transgenic mice and their non-transgenic litter-mates. Croton Oil was applied

directly to the ears of mice and DNFB was applied to the ears of mice after sensitization on shaved abdominal skin. Ear swelling was measured with a micrometer. In the ear-swelling test, the transgenic mice were divided into two groups; one group was treated with the murine monoclonal antibody ACK2, which blocks the interaction of SCF with its receptor (KIT), and the other group was treated with only saline. In addition, shaved abdominal skin of some mice was also treated with Croton Oil or with DNFB, and observed for inflammation and hyperpigmentation.

For statistical analysis, we used a mixed effects model, which allows us to fit repeated measurements over time and to compare different groups over time. We also performed orthogonal contrasts to evaluate the difference between treated and control groups at each time point. Immunoperoxidase study, using anti human SCF monoclonal antibodies, were performed in skin by standard method (7).

Experimental Results

Dermal mast cells accumulate in the presence of membrane/soluble keratinocyte SCF. Sections of skin from all mice producing membrane/soluble SCF (transgene one) showed increased mast cells in the dermis (Figure 2). In newborn transgene one positive mice, the mast cells were superficial near the dermal-epidermal junction, close to the epidermal source of soluble SCF (Figure 2a). In older mice the mast cells filled the papillary dermis in some areas, but were also present in the reticular dermis, in a pattern identical to that of human mastocytosis (Figure 2, b-d). Electron microscopic analysis confirmed the presence of numerous mast cells with characteristic granules within the dermis of the transgene one positive animals, and also showed that some of the heavily pigmented cells within the dermis of

transgene one positive mice were melanocytes (Figure 3). Mast cells were relatively rare and dermal melanocytes were not detected in the body wall skin of non-transgenic littermates and in transgene two positive animals of equivalent age. These observations were true across a wide range of copy numbers and levels of SCF mRNA expression (Figure 4). Since the keratin 14 promoter is properly expressed in the skin only by keratinocytes, and since the production of only membrane-bound keratinocyte SCF did not spontaneously result in increased dermal mast cells in transgene two positive animals, keratinocyte production of the soluble form of SCF appears to be able to cause cutaneous mastocytosis in mice.

SCF transgenic mice are hyperpigmented:

Targeted expression of each of the SCF transgenes in murine skin caused a similar, distinctive pigmented phenotype. The pigment responsible for the coat color of normal mice resides in the hair follicles and hair shafts, not in the epidermis. The transgenic mice, however, developed prominent epidermal pigmentation (Figure 5). Transgene positive animals could be identified by increased pigment at birth. By approximately 21 days of age, the phenotypes were well established; phenotype positive animals showed pigmentation of most of the skin as well as increased pigmentation of most of their skin as well as coat pigment. Extensive pigmentation was noted in a number of areas including the nose, mouth, ears, paws, and external genitalia when compared to normal littermate controls. There was enough individual variation in pigmentation so that no clear correlation between the level of pigmentation and the levels of transgenic expression could be shown. All transgenic animals showed similar degrees of pigmentation regardless of transgene type, copy number, or levels of SCF mRNA expression. In addition to the epidermal pigmentation, the three

transgene two positive agouti founders showed thin black transverse strips, consistent with the pigment distribution of the allophenic mice described by Beatrice Mintz (pictures not shown) (36).

5

Numerous melanocytes are maintained in the skin of transgenic mice:

10 The increased pigmentation of the skin of the transgene positive mice of both types is attributable to the presence of intraepidermal melanocytes, and to the epidermal melanin produced by those cells. Intraepidermal melanocytes can be identified in hematoxylin and eosin stained sections as cells in the basilar layers surrounded by clear halos (Figure 6, a & b) or in immunoperoxidase preparations by their expression of S-100 protein. Immunohistochemical analysis of animals expressing each of the transgenes showed numerous S-100(+) intraepidermal melanocytes (please also see Figure 2a). These melanocytes can be differentiated from Langerhans cells, which also express S-100 protein, because melanocytes are in the basal layers and Langerhans are in the suprabasal layers. Melanocytes can also be differentiated from Langerhans cells by their expression of the kit protein, the receptor for SCF, which is not expressed by Langerhans cells. Staining of transgenic animal skin with anti-kit antibody identified well-developed dendritic cells within the basilar layers of the epidermis and follicular epithelium, consistent with melanocytes (Figure 6c and 2b).

30

Histologic examination confirmed the presence of pigment within the epidermis of both transgene one and transgene two phenotype positive mice from all sites examined, including the ears, tail, footpads, and body wall (Figure 6a). In addition, transgene one positive mice showed many pigmented cells within the dermis. Pigmentary

35

abnormalities were not observed in transgene negative littermates. Only slight epidermal pigment was identified in these control mice, and mostly in non-hair bearing areas like the footpad and tail. Although pigment patterns were stable throughout much of the adult life of the mice, an occasional TG1 (msSCF) mouse developed patchy areas of depigmentation, mostly in the ears, associated with loss of epidermal melanocytes and increased pigment incontinence. This phenomenon was not observed in the mSCF mice.

Electron microscopy confirmed the presence of numerous melanocytes within the epidermis of both types of transgenic mice (Figure 7). Pigmented keratinocytes, similar to those seen in the epidermis of humans, were also present in the interadnexal epidermis of the transgenic mice. Intraepidermal melanocytes and pigmented keratinocytes were extremely rare in control mice.

All transgenic mice showed ear swelling which was greater in magnitude and more prolonged than the non-transgenic (B6) control mice:

Blocking SCF by administration of ACK2 decreased the magnitude of ear swelling in transgenic mice, as shown in the following figure 8.

Averaging across time:

There is a difference between the ACK2 treated and the control saline treated transgenic mice, which is significant at the 0.05 level. Averaging across time, there is also a significant difference between each of the two groups of transgenic mice (ACK2 treated and control) and the non-transgenic mice. Both comparisons are statistically significant at the 0.0001 level. See Figure 8. Observation of the abdominal wall skin treated with Croton Oil or DNFB showed hyperpigmentation and

thickening which was not observed in non-transgenic mice control (B6) mice that were treated identically. Histologically, hyperpigmentation correlated with dermal melanophages and increased epidermal melanin, identical to the changes seen in human postinflammatory hyperpigmentation.

Discussion

Melanocytes are maintained in human epidermis throughout life. In normal mice DOPA reaction positive cells (melanoblasts and melanocytes) are found in the epidermis at birth, but their number decreases from postnatal day 4 and is severely reduced after one month of age (37). One possible explanation for the maintenance of epidermal melanocytes in human skin, and the difference between the distribution of melanocytes in adult human and murine skin, could be expression of epidermal SCF. Human epidermal keratinocytes produce SCF (7, 8, 39), but the SCF gene does not appear to be expressed in murine epidermis (9). The results presented here show that SCF expression by murine epidermal keratinocytes causes the maintenance and stimulation of epidermal melanocytes throughout life. These data support the hypothesis that the decrease in melanocyte numbers in the postnatal mouse epidermis is due to a lack of local SCF expression. In combination with the fact that the soluble SCF produced by Sl/Sld mice is insufficient to support normal melanocyte survival and the observations that membrane-bound SCF promotes longer lasting kit activation and increased survival of kit dependent cells in the hematopoietic system (40,41), our data suggest that it is specifically the membrane-bound form of SCF that is crucial for melanocyte survival and function.

It is interesting to note that none of the animals expressing either of the transgenes described in this paper have developed melanoma to date, a finding which

supports previous observations that stimulation of the
kit tyrosine kinase receptor does not appear to promote
the development of melanocytic tumors (40). It also
seems likely that the animals described herein, or
5 animals derived from them, will be useful in the study of
cutaneous mastocytosis and epidermal melanocyte biology.

The fact that SCF transgenic mice have greater responses
to allergic and irritant contactants shows that epidermal
10 SCF can actively contribute to eczematous dermatitis.
This interpretation is confirmed by our demonstration
that the inflammation can be diminished by blocking the
SCF receptor with the ACK2 monoclonal antibody. Since
human post natal epidermal keratinocytes express SCF,
15 unlike post natal murine epidermal keratinocytes, and
alterations of human epidermal SCF are found in
spongiotic dermatitis (a form of eczema), these
observations also support our contention that the skin of
mice expressing epidermal SCF is a better model of human
20 skin than is the skin of normal mice. Further supporting
this claim is our previous observation of increased
soluble epidermal SCF in the hyperpigmented lesions of
mastocytosis. In sum, these data support our claim that
animals expressing epidermal SCF are more suitable for a
25 wide variety of investigations than those which do not.

REFERENCES

1. Silvers, W. K. (1979) "The coat colors of mice: a
model for mammalian gene action and interaction"
30 Springer-Verlag, Inc., New York. pp. 4-5 and
references therein;
2. Mayer, T. C. (1970) "A comparison of pigment cell
development in albino, steel, and dominant-spotting
35 mutant mouse embryos" Develop. Biol., 23:297-309;
3. Russell, E. S. (1979) "Hereditary anemias of the

mouse: a review for geneticists" Adv. Genet.
20:357-459;

- 5 4. Yarden, Y., et al. (1987) "Human proto-oncogene
 c-kit: a new cell surface receptor tyrosine kinase
 for an unidentified ligand" EMBO J., 6:3341-3351;
- 10 5. Qiu, F. H., et al. (1988) "Primary structure of
 c-kit: relationship with the CSF-1/PDGF receptor
 kinase family -- oncogenic activation of v-kit
 involves deletion of extracellular domain and C
 terminus" EMBO J., 7:1003-1011;
- 15 6. Geissler, E. N., et al. (1988) "The dominant-white
 spotting (W) locus of the mouse encodes the c-kit
 proto-oncogene" Cell, 55:185-192;
- 20 7. Longley, B. J. Jr, et al. (1993) "Altered metabolism
 of mast-cell growth factor (c-kit ligand) in
 cutaneous mastocytosis" N. Engl. J. Med.,
 328:1302-1307;
- 25 8. Weiss, R. R., et al. (1995) "Human dermal endothelial
 cells express membrane-associated mast cell growth
 factor" J. Invest. Dermatol., 104:101-106;
- 30 9. Yoshida, H., et al. (1996) "Neural and skin cell
 specific expression pattern conferred by Steel
 factor regulatory sequence in transgenic mice"
 Developmental Dynamics, 207:222-232;
- 35 10. Anderson, D. M., et al. (1990) "Molecular cloning of
 mast cell growth factor, a hematopoietin that is
 active in both membrane bound and soluble forms"
 Cell, 63:235-243;
11. Zsebo, K. M., et al. (1990) "Identification,
 purification, and biological characterization of

hematopoietic stem cell factor from Buffalo rat liver-conditioned medium" Cell, 63:195-201;

- 5 12. Flanagan, J. G. and P. Leder (1990) "The kit ligand: a cell surface molecule altered in steel mutant fibroblasts" Cell, 63:185-194;
- 10 13. Onoue, H., et al. (1989) "Suppressive effects of S1/Sld mouse embryo-derived fibroblast cell lines on diffusible factor-dependent proliferation of mast cells" Blood, 74:1557-1562;
- 15 14. Anderson, D. M., et al. (1991) "Alternate splicing of mRNAs encoding human mast cell growth factor and localization of the gene to chromosome 12q22-q24" Cell Growth & Development, 2:373-378;
- 20 15. Lu, H. S., et al. (1991) "Amino acid sequence and post-translational modification of stem cell factor isolated from buffalo rat liver cell-conditioned medium" J. Biol. Chem., 266:8102-8107;
- 25 16. Flanagan, J. G., et al. (1991) "Transmembrane form of the kit ligand growth factor is determined by alternative splicing and is missing Sld mutant" Cell, 64:1025-1035;
- 30 17. Brannan, C. I., et al. (1991) "Steel-Dickie mutation encodes a c-Kit ligand lacking transmembrane and cytoplasmic domains" Proc. Natl. Acad. Sci. USA, 88:4671-4674;
- 35 18. Zsebo, K. M., et al. (1990) "Stem cell factor is encoded at the S1 locus of the mouse and is the ligand for the c-kit tyrosine kinase receptor" Cell, 63:213-224;

19. Huang, E. J., et al. (1992) "Differential expression and processing of two cell associated forms of the kit-ligand: KL-1 and KL-2" Mol. Biol. Cell, 3:349-362;
- 5
20. Wehrle-Haller, B. and J. A. Weston (1995) "Soluble and cell-bound forms of steel factor activity play distinct roles in melanocyte precursor dispersal and survival on the lateral neural crest migration pathway" Development, 121:731-742;
- 10
21. Tsai, M., et al. (1991) "The rat c-kit ligand, stem cell factor, induces the development of connective tissue-type and mucosal mast cells in vivo: Analysis by anatomical distribution, histochemistry, and protease phenotype" J. Exp. Med., 174:125-131;
- 15
22. Harrist, T. J., et al. (1995) "Recombinant human stem cell factor (SCF) (c-kit ligand) promotes melanocyte hyperplasia and activation in vivo" Lab. Invest., 72:48A;
- 20
23. Costa, J. J., et al. (1996) "Recombinant human stem cell factor (KIT ligand) promotes human mast cell and melanocyte hyperplasia and functional activation in vivo" J. Exp. Med., 183:2681-2686;
- 25
24. Longley, B. J., et al. (1995) "The mast cell and mast cell disease" J. Am. Acad. Dermatol., 32:545-561;
- 30
25. Longley, B. J., et al. (1996) "Somatic c-KIT activating mutation in urticaria pigmentosa and aggressive mastocytosis: establishment of clonality in a human mast cell neoplasm" Nature Genetics, 12:312-314;
- 35
26. Furitsu, T., et al. (1993) "Identification of

mutations in the coding sequence of the proto-oncogene c-KIT in a human mast cell leukemia cell line causing ligand-independent activation of c-KIT product" J. Clin. Invest., 92:1736-1744;

5

27. Vassar, R., et al. (1989) "Tissue-specific and differentiation-specific expression of a human K14 keratin gene in transgenic mice" Proc. Natl. Acad. Sci. USA, 86:1563-1567;

10

28. Williams, D. E., et al. (1990) "Identification of a ligand for the c-kit proto-oncogene" Cell, 1990;63:167-174;

15

29. Majumdar, M. K., et al. (1994) "Identification and mutation of primary and secondary proteolytic cleavage sites in murine stem cell factor cDNA yields biologically active, cell-associated protein" J. Biol. Chem., 269:1237-1242;

20

30. Yasunaga, M., et al. (1995) "Cell cycle control of c-kit-1 IL-7R1B precursor cells by two distinct signals derived from IL-7 receptor and c-kit in a fully defined medium" J. Exp. Med., 182:315-323;

25

31. Kunisada, T., et al. (1996) "Characterization and isolation of melanocyte progenitors from mouse embryos" Development Growth & Differentiation, 38:87-97;

30

32. Yoshida, H., et al. (1996) "Distinct stages of melanocyte differentiation revealed by analysis of nonuniform pigmentation patterns" Development, 122:1207-1214;

35

33. Scott, J. E. and R. T. Mowry (1970) "Alcian blue - a consumer's guide" J. Histochem. Cytochem., 18:842;

34. Nishikawa, S., et al. (1991) "In utero manipulation of coat color formation by a monoclonal anti-c-kit antibody: two distinct waves of c-kit-dependency during melanocyte development" EMBO J., 10:2111-2118;
35. Okura, M., et al. (1995) "Effects of monoclonal anti-c-kit antibody (ACK2) on melanocytes in newborn mice" J. Invest. Dermatol., 105:322-328;
36. Bradl, M., et al. (1991) "Clonal coat color variation due to a transforming gene expressed in melanocytes of transgenic mice" Proc. Natl. Acad. Sci. USA, 88:6447-6451;
37. Grichnik, J. M., et al. (1995) "Human recombinant stem-cell factor induces melanocytic hyperplasia in susceptible patients" J. Am. Acad. Dermatol., 33:577-583;
38. Hirobe, T. (1984) "Histochemical survey of the distribution of the epidermal melanoblasts and melanocytes in the mouse during fetal and postnatal periods" Anat. Rec., 208:589-594;
39. Hamann, K., et al. (1995) "Expression of stem cell factor in cutaneous mastocytosis" Br. J. Dermatol., 133:203-208;
40. Funasaka, Y., et al. (1992) "C-kit-kinase induces a cascade of protein tyrosine phosphorylation in normal human melanocytes in response to mast cell growth factor and stimulates mitogen-activated protein kinase but is down-regulated in melanomas" Mol. Biol. Cell, 3:197-209.

SECOND SERIES

Mastocytosis is a neoplastic disease caused at least in part by somatic mutations of the *c-KIT* proto-oncogene resulting in constitutive activation of its protein product, KIT, the receptor tyrosine kinase for stem cell factor. KIT stimulates mast cell proliferation and prevents apoptosis of neoplastic mast cells. Human gastrointestinal stromal tumor cells also express mutated and activated kit (Hirota et al 1998). To develop potential therapies for mastocytosis and gastrointestinal stromal tumor cells we used indolinones, small molecules which inhibit tyrosine kinases.

The proto-oncogene *c-KIT* encodes KIT (Yarden et al, 1987; Qiu et al, 1988), the receptor tyrosine kinase for stem cell factor (Martin et al, 1990), also known as mast cell growth factor. Somatic *c-KIT* mutations causing ligand-independent activation of KIT and cell transformation (Furitsu et al, 1993; Kitayama et al, 1995; Tsujimura et al, 1996; Hirota et al, 1998; Ma et al, 1999a) appear causal in certain types of mastocytosis (Nagata et al, 1995; Longley et al, 1996, 1999; Ma et al, 1999a).

Documented activating *c-KIT* mutations fall into two groups. One group consists of mutations in codon 816 of human *c-KIT*, or its equivalent positions in other species, resulting in single residue substitution for Asp816 in the activation loop of the receptor kinase domain (Ma et al, 1999a). The other group of activating mutations includes single residue substitutions and in-frame insertions or deletions in the receptor intracellular juxtamembrane region, which disrupt intramolecular inhibition of the kinase by a putative juxtamembrane α -helix (Ma et al, 1999b). All sporadic

adult-onset mastocytosis patients examined to date, and
a subset of pediatric cases with atypical clinical
presentations, have activating codon 816 mutations
(Longley et al, 1999), whereas activating juxtamembrane
5 mutations are common in canine mastocytomas (Ma et al,
1999a) and in human gastrointestinal stromal tumors
(Hirota et al, 1998).

10 References for second series

Furitsu T, Tsujimura T, Tono T, Ikeda H, Kitayama H, et
al: Identification of mutations in the coding sequence
of the proto-oncogene *c-kit* in a human mast cell leukemia
cell line causing ligand-independent activation of *c-kit*
15 product. *J Clin Invest* 92: 1736-1744, 1993

Hirota S, Isozaki K, Moriyama Y, Hashimoto K, Nishida T,
et al: Gain-of-function mutations of *c-kit* in human
gastrointestinal stromal tumors. *Science* 279: 577-580,
20 1998

Kitayama H, Kanakura Y, Furitsu T, Tsujimura T, Oritani
K, et al: Constitutively activating mutations of *c-kit*
receptor tyrosine kinase confer factor-independent growth
and tumorigenicity of factor-dependent hematopoietic cell
25 lines. *Blood* 85: 790-798, 1995

Longley BJ, Tyrrell L, Lu S, Ma Y, Langley K, et al:
Somatic *c-KIT* activating mutation in urticaria pigmentosa
and aggressive mastocytosis: establishment of clonality
30 in a human mast cell neoplasm. *Nature Genet* 12: 312-314,
1996

Longley BJ, Metcalfe DD, Tharp M, Wang X, Tyrrell L, et

- al: Activating and dominant inactivating *c-KIT* catalytic domain mutations in distinct clinical forms of human mastocytosis. *Proc Natl Acad Sci USA* 96: 1609-1614, 1999
- 5 Ma Y, Longley BJ, Wang X, Blount JL, Langley K, Caughey GH: Clustering of activating mutations in *c-KIT*'s juxtamembrane coding region in canine mast cell neoplasms. *J Invest Dermatol* 112: 165-170, 1999a
- 10 Ma Y, Cunningham ME, Wang X, Ghosh I, Regan L, Longley BJ: Inhibition of spontaneous receptor phosphorylation by residues in a putative α -helix in the *KIT* intracellular juxtamembrane region. *J Biol Chem* 274: 13399-13402, 1999b
- 15 Martin FH, Suggs SV, Langley KE, Lu HS, Ting J, et al: Primary structure and functional expression of rat and human stem cell factor DNAs. *Cell* 63: 203-211, 1990
- 20 Nagata H, Worobec AS, Oh CK, Chowdhury BA, Tannenbaum S, et al: Identification of a point mutation in the catalytic domain of the protooncogene *c-kit* in peripheral blood mononuclear cells of patients who have mastocytosis with an associated hematologic disorder. *Proc Natl Acad Sci USA* 92: 10560-10564, 1995
- 25 Tsujimura T, Morimoto M, Hashimoto K, Moriyama Y, Kitayama H, et al: Constitutive activation of *c-kit* in FMA3 murine mastocytoma cells caused by deletion of seven amino acids at the juxtamembrane domain. *Blood* 87: 273-283, 1996
- 30 Yarden Y, Kuang W-J, Yang-Feng T, Coussens L, Munemitsu S, et al: Human proto-oncogene *c-kit*: a new cell surface

receptor tyrosine kinase for an unidentified ligand. *EMBO J* 6: 3341-3351, 1987

Third Series of Experiments

5 Passive Anaphylaxis: SCF and KIT stimulation have a number of effects on mast cells in vitro, but it is not clear what the overall of effects of blocking KIT would be on inflammation and physiology in vivo. For instance, numerous studies have shown that recombinant SCF causes
10 mast cell activation, directly stimulates mediator release, and alters the threshold for IgE dependent mediator release in vitro (Nakajima et al. Biochem. Biophys. Res. Commun. 1992) (Coleman et al. J. Immunol. 1993) (Columbo et al. 1992) (Bischoff et al. J. Exp. Med. 1992). Furthermore, a single injection of recombinant SCF causes mast cell activation in mice (Werschil et al. J. Exp. Med. 1992). However, daily administration of recombinant SCF to mice results in mast cell hyperplasia which varies at different anatomic sites, but does not
20 result in mast cell activation (Anado et al. J. Clin. Invest. 1993), and the chronic administration of recombinant SCF to mice has variable effects on passive anaphylaxis reactions. Chronic stimulation with recombinant SCF followed by elicitation of passive
25 anaphylaxis results in increased mast cell activation at some anatomic sites but not others and, surprisingly, a decrease in deaths in mice (Anado et al. J. Clin. Invest. 1993). This last result is counter-intuitive and indicates that exact effects of blocking KIT activation
30 in vivo cannot be predicted accurately based on in vitro data. The authors themselves describe the results as unexpected.

35 To investigate the effects of blocking KIT in vivo, the ACK2 KIT-blocking-antibody and passive cutaneous anaphylaxis were used in the following experiments:

The skin of Balb/c mice was sensitized by an intradermal injection of murine monoclonal IgE specific for dinitrophenyl hapten. A second set of mice was sensitized with both the IgE and ACK2. 24 hours later, dinitrophenylated human serum albumin was injected into the tail veins of the mice in a mixture of 1% Evans' Blue dye (1 mg). Mice were observed for anaphylaxis, and thirty minutes later were sacrificed. The dorsal skin at the area of sensitization was removed for measurement of the amount of extravasated dye. The amount of dye was determined colorometrically after extraction of the skin in 1 ml of 1 Normal KOH overnight, at 37 degrees centigrade. 0.6 Normal phosphoric acid in acetone (15:13) was added and the mixture cleared by centrifugation. The absorbent intensity (OD) of the supernatant was measured at 620nm spectrophotometrically. The amount of extravasated dye, which is a measure of anaphylaxis, was significantly reduced in the presence of ACK2 (mean OD 0.0223) when compared to animals sensitized without ACK2 (OD 0.0617) a difference which was significant with $p < 0.01$, and the animals treated with ACK2 showed less respiratory distress compared to animals without ACK2.

These data demonstrate, for the first time, that blocking the SCF-KIT signaling pathway in vivo can decrease anaphylaxis and bronchospasms. Combined with the data on cutaneous inflammation, they demonstrate that the SCF-KIT pathway is actively involved in inflammatory reactions in vivo, rather than playing merely a supportive role in the development and maintenance of KIT expressing cells. This critical finding shows that acute blockage of the SCF-KIT signaling pathway can inhibit inflammation.

References for Third Series of Experiments:

1. Nakajima K, Hirai K, Yamaguchi M, Takaishi T, Ohta K, Morita Y, Ito K. Biochem. Biophys. Res. Commun.

1992, 183:1076-83.

2. Coleman JW, Holliday MR, Kimber I, Zsebo KM and Galli SJ. J. Immunol. 1993, 150: 556-62.
- 5 3. Columbo M , Horowitz EM, Botana LM, MacGlashan DW Jr., Bochner BS, Gillis S, Zsebo KM, Galli SJ, Lichtenstein LM. J. Immunol. 1992, 149:599-608.
- 10 4. Bischoff SC and Dahinden CA. J. Exp. Med. 1992, 175:237-44.
5. Wershil BK, Tsai M, Geissler EN, Zsebo KM, Galli SJ. J. Exp. Med. 1992, 175:245-55.
- 15 6. Ando A, Martin TR, Galli SJ. J. Clin. Invest. 1993, 92:1639-49.

20

What is claimed:

- 5 1. A method of preventing or treating in a subject contact dermatitis which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat contact dermatitis so as to thereby prevent or treat contact dermatitis in the subject.
- 10 2. A method of preventing or treating in a subject hyperpigmentation which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat hyperpigmentation so as to thereby prevent or treat hyperpigmentation in the subject.
- 15 3. A method of preventing or treating in a subject asthma which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat asthma so as to thereby prevent or treat asthma in the subject.
- 20 4. A method of preventing or treating in a subject cutaneous inflammation which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent or treat cutaneous inflammation so as to thereby prevent or treat cutaneous inflammation in the subject.
- 25 5. A method of preventing or treating in a subject anaphylaxis and bronchospasm which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling
- 30
- 35

5 6. A method of preventing or treating in a subject
 mastocytosis which comprises administering to the
 subject an amount of a compound capable of
 inhibiting the stem cell factor signaling pathway
 effective to prevent or treat mastocytosis so as to
10 thereby prevent or treat mastocytosis in the
 subject.

20 8. A method of preventing or treating in a subject
hypersensitivity reactions which comprises
administering to the subject an amount of a compound
capable of inhibiting the stem cell factor signaling
pathway effective to prevent or treat urticaria so
25 as to thereby prevent or treat urticaria in the
subject.

10. The method of claim 9, wherein the tumor is a

gastrointestinal stromal tumor.

11. The method of claim 9, wherein the tumor is a germ cell tumor.

5

12. The method of any one of claims 1-9, which comprises inhibiting the kinase enzymatic reaction of kit protein.

10

13. The method of any one of claims 1-9, which comprises inhibiting chymase, elastase or other SCF cleaving enzymes.

15

14. The method of any one of claims 1-9, which comprises inhibiting ligand binding with an antibody, peptide, or nonpeptide chemical.

20

15. The method of any one of claims 1-9, which comprises inhibiting kit dimerization with an antibody, peptide, or nonpeptide chemical.

25

16. The method of any one of claims 1-9, wherein downstream signaling of the kit activation pathway is inhibited by blocking substrate association with kit kinase domain.

30

17. The method of any one of claims 1-9, wherein downstream signaling of the kit activation pathway is inhibited by blocking enzymatic function in the downstream signaling pathway.

35

18. The method of any one of claims 1-9, wherein downstream signaling of the kit activation pathway is inhibited by blocking binding of molecules in the downstream signaling pathway.

19. The method of any one of claims 1-9, wherein the

compound is an antibody or portion thereof.

- 5
20. The method of claim 19, wherein the antibody is a monoclonal antibody.
21. The method of claim 20, wherein the monoclonal antibody is a human, humanized or a chimeric antibody.
- 10 22. The method of claim 20, wherein the monoclonal antibody is an anti-kit antibody.
23. The method of claim 22, wherein the anti-kit antibody is ACK2.
- 15 24. The method of any one of claims 1-9, wherein the compound comprises a Fab fragment of an anti-kit antibody.
- 20 25. The method of any one of claims 1-9, wherein the compound comprises the variable domain of an anti-kit antibody.
- 25 26. The method of any one of claims 1-9, wherein the compound comprises one or more CDR portions of an anti-kit antibody.
- 30 27. The method of claim 19, wherein the antibody is selected from the group consisting of IgA, IgD, IgE, IgG and IgM.
- 35 28. The method of any one of claims 1-9, wherein the compound comprises a peptide, peptidomimetic, a nucleic acid, or an organic compound with a molecular weight less than 500 Daltons.
29. The method of any one of claims 1-9, wherein the

compound is sSCF, sKIT ligand or a fragment thereof.

30. The method of any one of claims 1-9, wherein the compound is sKIT or a fragment thereof.

5

31. The method of any one of claims 1-9, wherein the subject is a mammal.

10 32. The method of claim 31, wherein the mammal is a human being, dog or cat,

15 33. The method of any one of claims 1-9, wherein the administration is intralesional, intraperitoneal, intramuscular, subcutaneous, intravenous, liposome mediated delivery, transmucosal, intestinal, topical, nasal, oral, anal, ocular or otic delivery.

20 34. A method of providing contraception to a subject which comprises administering to the subject an amount of a compound capable of inhibiting the stem cell factor signaling pathway effective to prevent conception so as to thereby provide contraception to the subject.

25 35. The method of claim 34, wherein the subject is a male subject.

30 36. The method of claim 34, wherein the subject is a female subject.

37. The method of claim 34, which comprises inhibiting the kinase enzymatic reaction of kit protein.

35 38. The method of claim 34, which comprises inhibiting chymase, elastase or other SCF cleaving enzymes.

**METHODS FOR INHIBITING CUTANEOUS
INFLAMMATION AND HYPERPIGMENTATION**


Abstract of the Disclosure

5 This invention provides a method of preventing or
treating in a subject contact dermatitis which comprises
administering to the subject an amount of a compound
capable of inhibiting the stem cell factor signaling
10 pathway effective to prevent or treat contact dermatitis in
so as to thereby prevent or treat contact dermatitis in
the subject. This invention also provides a methods of
preventing or treating in a subject hyperpigmentation,
asthma, cutaneous inflammation, anaphylaxis and
15 bronchospasm, mastocytosis, tumors which express
activated kit, and conception.

20


25

FIG. 1A



TRANSGENE 1

FIG. 1B



TRANSGENE 2

FIG. 2A



FIG. 2B



FIG. 2C





FIG. 2D




FIG. 2E

FIG. 3A



FIG. 3B




FIG. 3C

FIG. 4






FIG. 5C

FIG. 6A




FIG. 6B



FIG. 6C




66622T 2444450

FIG. 7A



FIG. 7B



05474473 13593
666237 82442450

FIG. 8
ACK2 Effect on DNFB-Induced Ear Swelling
1X Mice 3/16/99




FIG. 9A





FIG. 9B



As a below-named inventor, I hereby declare that:

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

the specification of which:
(check one)

 was filed on as

Application Serial No _____

and was amended _____ (if applicable)

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56

Prior Foreign Application(s)

Priority Claimed

[illegible]

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No</u>	<u>Filing Date</u>	<u>Status</u>
N/A		

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No</u>	<u>Filing Date</u>	<u>Status</u>
09/306,143	May 6, 1999	Pending

And I hereby appoint

John P. White (Reg. No. 28,678); Christopher C. Dunham (Reg. No. 22,031); Norman H. Zivin (Reg. No. 25,385); Jay H. Maioli (Reg. No. 27, 213); William E. Pelton (Reg. No. 25, 702); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Wendy E. Miller (Reg. No. 35,615); Richard S. Milner (Reg. No. 33, 970); Robert T. Maldonado (Reg. No. 38,232); Paul Teng (40,837); Gary J. Gershik (Reg. No. 39,992); Richard F. Jaworski (Reg. No. 33, 515); Elizabeth M. Wiechowski (Reg. No. 42, 226); and Pedro C. Fernandez (Reg. No. 41, 741)

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

Please address all communications, and direct all telephone calls, regarding this application to:

John P. White _____ Reg No. 28,678
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
Tel. (212) 278-0400

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or
first joint inventor B. Jack Longley _____

Inventor's signature _____

Citizenship United States of America Date of signature _____

Residence 49 Woodstock Road, Hamden, CT 06517 _____

Post Office Address same as above _____

Full name of joint
inventor (if any) _____

Inventor's signature _____

Citizenship _____ Date of signature _____

Residence _____

Post Office Address _____

Full name of joint
inventor (if any) _____

Inventor's signature _____

Citizenship _____ Date of signature _____

Residence _____

Post Office Address _____
